

TABLE 7.—Estimated relative risks for current and former cigarette smokers, females aged 35 years or more, 4-year (1982–86) followup of American Cancer Society 50-State study (CPS-II)

Underlying cause of death	Current smokers ^a	Former smokers ^a
All causes	1.90 (1.82–1.98) ^b	1.32 (1.27–1.37) ^b
CHD, age ≥35 (410–414) ^c	1.78 (1.62–1.97)	1.31 (1.19–1.44)
CHD, age 35–64 ^d (410–414)	3.00 (2.50–3.59)	1.43 (1.15–1.77)
CHD, age ≥65 (410–414)	1.60 (1.42–1.80)	1.29 (1.16–1.43)
Other Heart Disease ^e (390–398, 401–405, 415–417, 420–429)	1.69 (1.44–1.99)	1.16 (1.00–1.34)
Cerebrovascular Lesions, age ≥35 (430–438)	1.84 (1.56–2.16)	1.06 (0.88–1.27)
Cerebrovascular Lesions, age 35–64 (430–438)	4.80 (3.52–6.54)	1.41 (0.94–2.13)
Cerebrovascular Lesions, age ≥65 (430–438)	1.47 (1.19–1.81)	1.01 (0.83–1.24)
Other Circulatory Disease ^f (440–448)	3.00 (2.20–4.08)	1.34 (0.95–1.90)
COPD (490–492, 496)	10.47 (7.78–14.09)	7.04 (5.33–9.30)
Other Respiratory Disease ^g (010–012, 480–489, 493)	2.18 (1.60–2.97)	1.38 (1.04–1.84)
Cancer, Lip, Oral Cavity, Pharynx (140–149)	5.59 (3.15–9.91)	2.88 (1.57–5.26)
Cancer, Esophagus (150)	10.25 (4.94–21.27)	3.16 (1.45–6.85)
Cancer, Pancreas (157)	2.33 (1.77–3.08)	1.78 (1.37–2.30)
Cancer, Larynx (161)	17.78 (3.45–91.74)	11.88 (2.46–57.34)
Cancer, Lung (162)	11.94 (9.99–14.26)	4.69 (3.86–5.70)
Cancer, Cervix Uteri (180)	2.14 (1.06–4.30)	1.94 (0.97–3.87)
Cancer, Kidney (189)	1.41 (0.86–2.30)	1.16 (0.72–1.87)
Cancer, Bladder, Other Urinary Organs (188)	2.58 (1.31–5.08)	1.85 (1.00–3.42)

NOTE: Preliminary estimates, based upon 2,418,909 woman-years of exposure among female subjects who never smoked regularly, or who smoked only cigarettes, present or past. Relative risks, estimated with respect to women who never smoked regularly, have been directly standardized to the age distribution of all woman-years of exposure.

^aRefers to cigarette smoking status at enrollment (September 1982).

^bNumbers in parentheses are 95-percent confidence intervals, computed on the assumption that the logarithm of relative risk was normally distributed.

^cAll disease codes refer to International Classification of Diseases, Ninth Revision.

^dWhen an age range is given, it refers to the age at enrollment in 1982.

^eIncludes Hypertensive Heart Disease (401–404).

^fIncludes Aortic Aneurysm, Non-Syphilitic, and General Arteriosclerosis (440–441).

^gIncludes Influenza and Pneumonia (480–487).

SOURCE: Unpublished tabulations, American Cancer Society.

tween cigarette use and risk of stroke. They also noted a slight increase in risk among former cigarette smokers, especially for the first 2 years after cessation. The preliminary results from CPS-II, reported in Tables 6 and 7, further support a causal role for cigarette smoking in stroke.

The preliminary results of CPS-II also show significantly higher relative risks for cancers of the lip, oral cavity and pharynx, esophagus, and lung, as compared with CPS-I. The computed relative risk for lung cancer death has increased to 22 in men and 12 in women. While the relative risks for COPD death have not changed significantly among men, there is a trend toward increasing risk among women. The available data from CPS-II do not permit identification of specific mortality risks for hypertensive heart disease, aortic aneurysm, and influenza and pneumonia, as in CPS-I. However, among broader categories of cardiovascular and nonneoplastic respiratory disease, increased risks are likewise found in CPS-II.

Endocrine and Sex-Related Cancers in Women

A protective effect of smoking on cancer of the endometrium has been suggested in a recent case-control study (Lesko et al. 1985). For CPS-I, the relative risk for cancers of the uterine corpus (ICD-7 Codes 172-174) among current smokers was 0.94 (95-percent confidence interval, 0.57 to 1.53). Preliminary results for CPS-II suggest a reduced relative risk for endometrial cancer (ICD-9 Code 182).

Recent data on a possible protective effect of smoking for breast cancer have been contradictory (See Chapter 2; Rosenberg et al. 1984). For CPS-I, the relative risk for breast cancer (ICD-7 Code 170) among current smokers was 0.88 (95-percent confidence interval, 0.77 to 1.01), while the relative risk among former smokers was 1.20 (95-percent confidence interval, 1.15 to 1.35). Preliminary data from CPS-II have likewise been contradictory.

An increased risk of cervical cancer among cigarette smokers has been reported in case-control studies (LaVecchia et al. 1986; Nischan, Ebeling, Schindler 1988). For CPS-I, the relative risk for cervical cancer (ICD-7 Code 171) was 1.10 (95-percent confidence interval, 0.83 to 1.47). Data from CPS-II show a twofold increase in cervical cancer mortality among current smokers (relative risk 2.14, 95-percent confidence interval 1.06 to 4.30).

Summary

The relative risks for current smokers for selected comparable disease categories causally related to smoking in CPS-I and CPS-II are summarized and listed side by side in Table 8. These comparisons show substantial increases in the risk of death due to smoking for most of the disease categories listed between the years 1959 and 1965 and 1982 and 1986. Statistically significant increases in relative risks occurred in those disease categories for which 95-percent confidence limits around the estimated relative risks do not overlap between CPS-I and CPS-II. Compared with men during this period, women experienced greater increases in the relative risks of cerebrovascular lesions (ages 35 to 64 years), COPD, laryngeal cancer, and lung cancer.

TABLE 8.—Summary of estimated relative risks for current cigarette smokers, major disease categories causally related to cigarettes, males and females aged 35 years and older, CPS-I (1959–65) and CPS-II (1982–86)

Underlying cause of death ^a	Males		Females	
	CPS-I	CPS-II	CPS-I	CPS-II
CHD, age ≥35	1.83	1.94	1.40	1.78 ^b
CHD, age 35–64	2.25	2.81 ^b	1.81	3.00 ^b
Cerebrovascular Lesions, age ≥35	1.37	2.24 ^b	1.19	1.84 ^b
Cerebrovascular Lesions, age 35–64	1.79	3.67 ^b	1.92	4.80 ^b
COPD	8.81	9.65	5.89	10.47
Cancer, Lip, Oral Cavity, and Pharynx	6.33	27.48	1.96	5.59
Cancer, Esophagus	3.62	7.60	1.94	10.25 ^b
Cancer, Pancreas	2.34	2.14	1.39	2.33
Cancer, Larynx	10.00	10.48	3.81	17.78
Cancer, Lung	11.35	22.36 ^b	2.69	11.94 ^b

^aSee Tables 4–7 for International Classification of Disease codes.

^b95-percent confidence intervals do not overlap between CPS-I and CPS-II.

SOURCE: Tables 4–7.

Smoking-Attributable Mortality in the United States, 1965 and 1985

Table 9 reports the attributable risks *a* from cigarette smoking during the year 1965. Ten causes of death are considered: CHD, COPD, cerebrovascular disease, and cancers of seven sites. The computations are based upon the age-adjusted relative risks reported in CPS-I and the prevalence rates reported in the 1965 NHIS. For men, the age-adjusted relative risks among present and past cigarette smokers with a history of pipe or cigar use were slightly lower than those for present and past smokers of cigarettes exclusively. While the latter are reported for comparison in Table 4, the former were used in the attributable risk computations. In 1965, as shown in Figure 2, about two-thirds of men with a history of regular cigarette smoking were also exposed to pipe or cigar smoke. (As noted in Note b of Table 10 below, the use of relative risks derived from the death rates of men who smoked cigarettes exclusively resulted in about a 5-percent increase in attributable deaths for 1965.) For women, the computation of attributable risks in 1965 did not distinguish between current and former smokers.

TABLE 9.—Estimated attributable risks for 10 selected causes of death from cigarette smoking, males and females, United States, 1965

Cause of death	Males ^a (%)	Females ^b (%)
CHD, age 35–64	42 (40–45) ^c	26 (23–30)
CHD, age ≥65	11 (9–14)	3.3 (2.1–5.1)
COPD	84 (79–88)	67 (57–76)
Cancer of lip, oral cavity, and pharynx	74 (59–85)	27 (12–51)
Cancer of larynx	84 (61–94)	47 (8–90)
Cancer of esophagus	57 (36–76)	14 (6–29)
Cancer of lung	86 (82–88)	40 (31–50)
Cancer of pancreas	41 (30–53)	14 (6–30)
Cancer of bladder	53 (39–66)	36 (20–56)
Cancer of kidney	36 (19–56)	17 (5–42)
Cerebrovascular disease, age 35–64	28 (21–36)	28 (22–33)
Cerebrovascular disease, age ≥65	2.0 (0.6–6.6)	1.3 (0.2–6.5)

^aFor males, computations based on prevalence rates in Table 2 and relative risks for male current and former cigarette smokers, with or without a history of pipe and cigar smoking, derived from CPS-1.

^bFor females, attributable risks computed from prevalence rates in Table 2 and relative risks for all female smokers, past and present, in Table 5.

^cNumbers in parentheses are 95-percent confidence intervals.

In 1965, as Table 9 reveals, cigarette smoking was responsible for 42 percent of CHD deaths among younger men and 26 percent of deaths among younger women. For COPD deaths at all ages, the smoking-attributable risks were 84 percent for men and 67 percent for women. For lung cancer, the respective attributable risks were 86 percent and 40 percent for men and women. With the exception of deaths from stroke among younger persons, attributable risks were markedly higher for men.

Table 10 reports the corresponding smoking-attributable deaths, *A*, during the year 1965. Attributable deaths were computed by multiplying the attributable risk percentages in Table 9 by the corresponding cause-specific death rates among persons aged 20

TABLE 10.—Estimated deaths (in thousands) attributable to cigarette smoking, 10 selected causes, males and females, United States, 1965

Cause of death	Males	Females
CHD, age <65	51 (48–54) ^a	9.5 (8.2–10.8)
CHD, age ≥65	25 (20–30)	6.0 (3.9–9.4)
COPD	16 (15–17)	2.3 (2.0–2.7)
Cancer of lip, oral cavity, and pharynx	3.6 (2.9–4.2)	0.4 (0.2–0.8)
Cancer of larynx	1.9 (1.4–2.2)	0.1 (0.02–0.3)
Cancer of esophagus	2.4 (1.5–3.2)	0.1 (0.2–0.8)
Cancer of lung	35 (34–36)	3.1 (2.4–3.8)
Cancer of pancreas	3.8 (2.8–4.9)	0.9 (0.4–2.0)
Cancer of bladder	3.0 (2.2–3.7)	1.0 (0.5–1.5)
Cancer of kidney	1.2 (0.7–1.9)	0.3 (0.1–1.8)
Cerebrovascular disease, age <65	5.5 (4.2–7.2)	4.7 (3.8–5.6)
Cerebrovascular disease, age ≥65	1.5 (0.4–4.8)	1.0 (0.2–5.9)
Ten causes	150 ^b (143–157)	30 (26–34)

NOTE: Computed from Table 9 and tabulations of deaths at ages 20 years or more by cause for 1965 (NCHS 1967).

Sums may not equal totals because of rounding.

^aNumbers in parentheses are 95-percent confidence intervals.

^bWhen the attributable risk estimates given in Note a of Table 9 were used, the total attributable deaths for males were 158,000 (95-percent confidence interval, 151,000 to 166,000). Approximately two-thirds of the 8,000 additional deaths were from CHD.

years or more. For the 10 causes combined, cigarette smoking was responsible for 150,000 deaths among men and 30,000 deaths among women in 1965.

Among men, CHD deaths made up 51 percent of smoking-attributable mortality for the 10 causes combined. This proportion is consistent with the estimate of 45 percent reported by the 1964 Advisory Committee to the Surgeon General for excess mortality from all causes (US PHS 1964). Similarly, lung cancer accounted for 23 percent of the smoking-attributable mortality for the 10 causes combined—again consistent with the

1964 Report's estimate of 16 percent of deaths from all causes. Among women, CHD deaths made up 52 percent and lung cancer 10 percent of the smoking-attributable mortality from the 10 causes combined.

Table 11 shows the estimated attributable risks *a* from cigarette smoking for the year 1985. For comparability with the 1965 calculations, the same 10 causes of death are considered. The computations are based upon the relative risks reported in CPS-II and the prevalence rates reported in the 1985 NHIS. For men, the computations employed the relative risks for past and present smokers of cigarettes exclusively, as shown in Table 6. As Figure 2 indicates, the proportion of male smokers who used other forms

TABLE 11.—Estimated attributable risks for 10 selected causes of death from cigarette smoking, males and females, United States, 1985

Cause of death	Males (%)	Females (%)
CHD, age <65	45 (40–50) ^a	41 (34–48)
CHD, age ≥65	21 (17–26)	12 (9–15)
COPD	84 (78–88)	79 (73–83)
Cancer of lip, oral cavity, and pharynx	92 (79–97)	61 (45–76)
Cancer of larynx	81 (57–93)	87 (56–97)
Cancer of esophagus	78 (62–89)	75 (57–87)
Cancer of lung	90 (88–92)	79 (75–82)
Cancer of pancreas	29 (18–43)	34 (25–44)
Cancer of bladder	47 (31–63)	37 (18–61)
Cancer of kidney	48 (32–64)	12 (3–43)
Cerebrovascular disease, age <65	51 (36–65)	55 (45–65)
Cerebrovascular disease, age ≥65	24 (16–35)	6 (2–14)

NOTE: Computed from Tables 2, 6, and 7. For adult men under 65, the proportions of current and former cigarette smokers in 1985 were, respectively, 34.7 and 25.8 percent. For men 65 or older, the prevalences of current and former cigarette smoking were, respectively, 19.4 and 51.1 percent. For adult women under 65, the corresponding proportions were 30.1 and 16.5 percent; for adult women 65 or older, 12.6 and 19.6 percent.

^aNumbers in parentheses are 95-percent confidence intervals.

of tobacco was too small to affect significantly the results for 1985. For women, relative risks for current and former cigarette smokers were employed (Table 7).

Comparison of Tables 9 and 11 reveals significant increases in attributable risk from 1965–85. In 1985, smoking accounted for 21 percent of CHD deaths in older men, compared with 11 percent in 1965. The attributable risks for cancers of the lip, oral cavity and pharynx, esophagus, and lung increased significantly.

Changes in the attributable risk estimates for women are even more striking. Among younger women, smoking now accounts for an estimated 41 percent of CHD deaths and an estimated 55 percent of lethal strokes, compared with 26 and 28 percent, respectively, in 1965. Among women of all ages, 79 percent of lung cancers are attributable to cigarette use (see Table 11).

Overall, smoking accounted for 86.7 percent of all lung cancer deaths (95-percent confidence interval 84.9 to 88.4), 81.8 percent of all COPD deaths (95-percent confidence interval 78.3 to 85.3), and 21.5 percent of all CHD deaths (95-percent confidence interval 19.4 to 23.4). In addition, smoking accounted for 18.0 percent of all stroke deaths (95-percent confidence interval 14.2 to 22.9).

Table 12 reports estimated smoking-attributable deaths for the 10 causes during 1985. Total deaths have increased to 231,000 for men and 106,000 for women. As opposed to 1965, CHD in men now accounts for only one-third of the smoking-attributable mortality from the 10 causes combined. The proportion of these attributable deaths due to lung cancer has increased to one-third. Likewise, among women, smoking-attributable CHD fatalities now account for one-third of the 10-cause total; the relative importance of smoking-induced cancer fatalities has also increased.

The total 10-cause smoking-attributable mortality for 1985 was 337,000 deaths, compared with 183,000 in 1965. A portion of the observed 1965–85 increase, however, was the result of population growth. In addition, there were increases in the proportion of elderly persons who would be more at risk for smoking-induced death. For men and women, respectively, Figures 10 and 11 show the results of a correction for population increase and population aging. In each figure, three quantities are shown for each of four categories of smoking-attributable mortality: CHD deaths under age 65; CHD deaths age 65 years or more; COPD deaths; and lung cancer deaths. The first quantity is the estimated smoking-attributable deaths for 1965. The second bar shows smoking-attributable deaths for 1985. The third bar shows the estimated 1985 smoking-attributable deaths if the U.S. populations at each age had remained at 1965 levels. The latter quantities were computed as aD^* , where a is the attributable risk given in Table 11 and D^* is a population-corrected estimate of 1985 U.S. deaths. The latter quantity was computed by multiplying 1985 age-specific death rates by the populations at risk in 1965.

Figures 10 and 11 show that population growth and aging cannot explain the changes in smoking-attributable mortality between 1965 and 1985. In particular, the marked increases in smoking-attributable deaths from lung cancer and COPD in women are systematic consequences of the American woman's adoption of lifelong cigarette smoking, from teenage years onward.

For men, population-corrected deaths due to smoking in 1985 were 165,000, compared with 150,000 in 1965. For women, population-corrected deaths due to smoking

TABLE 12.—Estimated deaths (in thousands) attributable to cigarette smoking, 10 selected causes, males and females, United States, 1985

Cause of death	Males	Females
CHD, age <65	34 (30–38) ^a	11 (9–12)
CHD, age ≥65	44 (36–54)	26 (20–34)
COPD	37 (35–39)	20 (18–21)
Cancer of lip, oral cavity, and pharynx	5.1 (4.4–5.4)	1.6 (1.2–2.0)
Cancer of larynx	2.3 (1.6–2.7)	0.6 (0.4–0.7)
Cancer of esophagus	5.0 (4.0–5.7)	1.6 (1.3–1.9)
Cancer of lung	76 (74–77)	30 (29–32)
Cancer of pancreas	3.3 (2.1–5.0)	3.4 (2.8–5.1)
Cancer of bladder	3.1 (2.1–4.2)	1.1 (0.6–1.9)
Cancer of kidney	2.6 (1.8–3.5)	0.4 (0.1–1.5)
Cerebrovascular disease, age <65	5.5 (3.9–7.0)	5.2 (4.3–6.2)
Cerebrovascular disease, age ≥65	12 (8–17)	4.8 (1.9–11.4)
Ten causes	231 (220–242)	106 (98–115)

NOTE: Computed from Table 11 and unpublished tabulations of deaths at ages 20 years or more by cause from NCHS, 1985. Sum of individual causes may not equal totals because of rounding.

^aNumbers in parentheses are 95-percent confidence intervals.

in 1985 were 67,000, compared with 30,000 in 1965. Even if the population had remained entirely stable during 1965 through 1985, the lethality of cigarette use in American women would have doubled.

Among men, the total of 231,000 smoking-induced deaths in 1985 represented 41 percent of total deaths from the 10 causes combined and 22 percent of all deaths among persons aged 20 years or more. Among women, the total of 106,000 smoking-induced deaths represented 25 percent of deaths from the 10 causes combined and 11 percent of deaths from all deaths among persons aged 20 years or more.

The computations in Tables 10 and 12 have omitted other causes of death that are likely to be attributable to cigarette use. If the relative risks given in Tables 6 and 7 for




FIGURE 10.—Estimated cigarette-smoking-attributable deaths from CHD, COPD, and lung cancer, males aged 20 years or more, United States, 1965 and 1985

NOTE: For the bars marked 1985*, the estimated smoking-attributable deaths in 1985 have been corrected for population increases during 1965–85.




FIGURE 11.—Estimated cigarette-smoking-attributable deaths from CHD, COPD, and lung cancer, females aged 20 years or more, United States, 1965 and 1985

NOTE: For the bars marked 1985*, the estimated smoking-attributable deaths in 1985 have been corrected for population increases during 1965–85.

the broader categories of cardiovascular and nonneoplastic respiratory disease are applied to deaths from hypertensive heart disease, arteriosclerosis, aortic aneurysm, and influenza and pneumonia, then smoking-attributable deaths would increase to 256,000 among men and 126,000 among women. Inclusion of deaths among newborns and infants due to smoking during pregnancy would add an additional 2,500 to the total (CDC 1987b; McIntosh 1984; Kleinman et al. 1988); this does not include fetal loss due to smoking (Stein et al. 1981). Inclusion of lung cancer deaths among nonsmokers due to environmental tobacco smoke (NRC 1986) would add 3,800 and inclusion of deaths from cigarette-caused fires (Hall 1987) would add 1,700 to total attributable deaths. Inclusion of deaths due to cervical cancer caused by smoking would add 1,500. Including these additional causes of death, the smoking-attributable mortality in 1985 is then estimated to be approximately 390,000. Recent studies have also noted increased risks among smokers for hepatic cancer (Trichopoulos et al. 1987), penile cancer (Hellberg et al. 1987), leukemia (Kinlen and Rogot 1988), and anal cancer (Daling et al. 1987).

Among all persons at risk during 1985, an estimated 52 million were also cigarette smokers in 1965. The remaining 42 million were new cigarette smokers. In 1985, only about 4,400 deaths occurred among the latter group, which consists of persons in their teens, twenties, and thirties. Thus, 99 percent of deaths attributable to cigarette use in 1985 occurred among people who started smoking in 1965 or earlier. The vast majority of these people started smoking before the release of the 1964 Surgeon General's Report

TABLE 13.—Estimated risks of various activities

Activity or cause	Annual fatalities per 1 million exposed persons
Active smoking	7,000 ^a
Alcohol	541
Accident	275
Disease	266
Motor vehicles	187
Alcohol-involved	95
Non-alcohol-involved	92
Work	113
Swimming	22
Passive smoking ^b	19
All other air pollutants ^b	6
Football	6
Electrocution	2
Lightning	0.5
DES in cattlefeed	0.3
Bee sting	0.2
Basketball	0.02

NOTE: Activities are not mutually exclusive; there are overlaps between categories. Differences in fatalities do not imply proportionate differences in years of life lost.

^aNumber of deaths per million smokers who began smoking before 1965.

^bCancer deaths only.

SOURCE: Active smoking, CPS-II; NHISs 1965, 1985; U.S. Bureau of the Census (1974, 1986). Other activities or causes, U.S. President (1987).

and before the 1965 Federal Cigarette Labeling and Advertising Act. For this group, the annual smoking-attributable fatality rate is about 7 deaths per 1,000 at risk, or about 7,000 deaths per 1 million persons. As shown in the Economic Report of the President (U.S. President 1987), this rate far exceeds the rates for other risks of death (Table 13).

Conclusions

1. Lung cancer death rates increased two- to fourfold among older male smokers over the two decades between the American Cancer Society's two Cancer Prevention Studies (CPS-I, 1959–65, and CPS-II, 1982–86). Lung cancer death rates for younger male smokers fell about 30 to 40 percent during this period.
2. Lung cancer death rates increased four- to sevenfold among female smokers aged 45 years or older in CPS-II compared with CPS-I, while lung cancer death rates among younger women declined 35 to 55 percent.
3. The two-decade interval witnessed a two- to threefold increase in death rates from chronic obstructive pulmonary disease (COPD) in female smokers aged 55 years or older.
4. There was no change in the age-adjusted death rates for lung cancer and COPD between CPS-I and CPS-II among men and women who never smoked regularly.
5. Overall death rates from coronary heart disease (CHD) declined substantially between CPS-I and CPS-II. The decline in CHD mortality among nonsmokers, however, was notably greater than among current cigarette smokers.
6. In CPS-II, the relative risks of death from cerebrovascular lesions were 3.7 and 4.8 for men and women smokers under age 65. Increased risks of stroke were also observed among older smokers and former smokers. Along with the recently reported results of other studies, these findings strongly support a causal role for cigarette smoking in thromboembolic and hemorrhagic stroke.
7. In 1985, smoking accounted for 87 percent of lung cancer deaths, 82 percent of COPD deaths, 21 percent of CHD deaths, and 18 percent of stroke deaths. Among men and women less than 65 years of age, smoking accounted for more than 40 percent of CHD deaths.
8. The large increase in smoking-attributable mortality among American women between 1965 and 1985 was a direct consequence of their adoption of lifelong cigarette smoking, especially from their teenage years onward.
9. In 1985, 99 percent of smoking-attributable deaths occurred among people who started smoking before the 1964 Surgeon General's Report. For this group, the annual smoking-attributable fatality rate is about 7,000 deaths per 1 million persons at risk.
10. For 10 causes of death, a total of 337,000 deaths were attributable to smoking in 1985. These represented 22 percent of all deaths among men and 11 percent among women. If other cardiovascular, neoplastic, and respiratory causes of death were included—as well as deaths among newborns and infants resulting from maternal smoking, deaths from cigarette-caused residential fires, and lung cancer deaths among nonsmokers due to environmental tobacco smoke—the total smoking-attributable mortality was about 390,000 in 1985.

References

- ABBOTT, R.D., YIN, Y., REED, D.M., YANO, K. Risk of stroke in male cigarette smokers. *New England Journal of Medicine* 315(12):717-720, September 18, 1986.
- ALDERSON, M.R., LEE, P.N., WANG, R. Risks of lung cancer, chronic bronchitis, ischaemic heart disease, and stroke in relation to type of cigarette smoked. *Journal of Epidemiology and Community Health* 39(4):286-293, December 1985.
- AMERICAN CANCER SOCIETY. Unpublished tabulations.
- ANDERSON, S., AUQUIER, A., HAUCK, W.W., OAKES, D., VANDAELE, W., WEISBERG, H.I. *Statistical Methods for Comparative Studies*. New York: John Wiley and Sons, 1980.
- BLACK, B., LILIENFELD, D.E. Epidemiological proof in toxic tort litigation. *Fordham Law Review* 52:732-785, 1984.
- BOND, V.P. The cancer risk attributable to radiation exposure: Some practical problems. *Health Physics* 40:108-111, January 1981.
- BOTKIN, J.R. The fire-safe cigarette. *Journal of the American Medical Association* 260(2):226-229, July 8, 1988.
- BRACKBILL, R., FRAZIER, T., SHILLING, S. Smoking characteristics of US workers, 1978-1980. *American Journal of Industrial Medicine* 13:5-41, 1988.
- BROWN, C.C., KESSLER, L.G. Projections of lung cancer mortality in the United States: 1985-2025. *Journal of the National Cancer Institute* 80(1):43-51, March 2, 1988.
- CASTELLI, W.P., DAWBER, T.R., FEINLEIB, M., GARRISON, R.J., MCNAMARA, P.M., KANNEL, W.B. The filter cigarette and coronary heart disease: The Framingham Study. *Lancet* 2(8238):109-113, July 18, 1981.
- CENTERS FOR DISEASE CONTROL. Cigarette smoking in the United States, 1986. *Morbidity and Mortality Weekly Report* 36(35):581-585, September 11, 1987a.
- CENTERS FOR DISEASE CONTROL. Smoking-attributable mortality and years of potential life lost—United States, 1984. *Morbidity and Mortality Weekly Report* 36(42):693-697, October 30, 1987b.
- COLDITZ, G.A., BONITA, R., STAMPFER, M.J., WILLETT, W.C., ROSNER, B., SPEIZER, F.E., HENNEKENS, C.H. Cigarette smoking and risk of stroke in middle-aged women. *New England Journal of Medicine* 318(15):937-941, April 14, 1988.
- CONSUMER PRODUCT SAFETY COMMISSION. *Toward a Less Fire-Prone Cigarette: Final Report of the Technical Study Group on Cigarette and Little Cigar Fire Safety*. Consumer Product Safety Commission, October 1987.
- CORNFIELD, J. A method of estimating comparative rates from clinical data. Applications to cancer of the lung, breast, and cervix. *Journal of the National Cancer Institute* 11:1269-1275, 1951.
- COX, L.A. JR. Statistical issues in the estimation of assigned shares for carcinogenesis liability. *Risk Analysis* 7(1):71-80, 1987.
- CURRENT POPULATION SURVEY. Unpublished tabulations, 1985.
- DALING, J.R., WEISS, N.S., HISLOP, T.G., MADEN, C., COATES, R.J., SHERMAN, K.J., ASHLEY, R.L., BEAGRIE, M., RYAN, J.A., COREY, L. Sexual practices, sexually transmitted diseases, and the incidence of anal cancer. *New England Journal of Medicine* 317(16):973-977, October 15, 1987.
- DEANFIELD, J., WRIGHT, C., KRIKLER, S., RIBEIRO, P., FOX, K. Cigarette smoking and the treatment of angina with propranolol, atenolol, and nifedipine. *New England Journal of Medicine* 310(15):951-954, April 12, 1984.

- DOLL, R., GRAY, R., HAFNER, B., PETO, R. Mortality in relation to smoking: 22 years' observations on female British doctors. *British Medical Journal* 280(6219):967-971, April 5, 1980.
- DOLL, R., HILL, A.B. Lung cancer and other causes of death in relation to smoking. A second report on the mortality of British doctors. *British Medical Journal* 2:1071-1081, November 10, 1956.
- DOLL, R., HILL, A.B. Mortality in relation to smoking: Ten years' observations of British doctors. *British Medical Journal* 1:1399-1410, May 30, 1964; 1460-1467, June 6, 1964.
- DOLL, R., HILL, A.B. Mortality of British doctors in relation to smoking: Observations on coronary thrombosis. In: Haenszel, W. (ed.) *Epidemiological Approaches to the Study of Cancer and Other Chronic Diseases*, NCI Monograph 19. U.S. Department of Health, Education, and Welfare, Public Health Service, National Cancer Institute, January 1966, pp. 205-268.
- DOLL, R., PETO, R. Mortality in relation to smoking: 20 years' observations on male British doctors. *British Medical Journal* 2:1525-1536, December 25, 1976.
- DOLL, R., PETO, R. Cigarette smoking and bronchial carcinoma: Dose and time relationships among regular and lifelong non-smokers. *Journal of Epidemiology and Community Health* 32(4):302-313, December 1978.
- DOLL, R., PETO, R. The causes of cancer: Quantitative estimates of avoidable risks of cancer in the United States today. *Journal of the National Cancer Institute* 66(6):1191-1308, June 1981.
- DORN, H.F. Tobacco consumption and mortality from cancer and other diseases. *Public Health Reports* 74(7):581-593, July 1959.
- GALAN, K.M., DELIGONUL, U., KERN, M.J., CHAITMAN, B.R., VANDORMAEL, M.G. Increased frequency of restenosis in patients continuing to smoke cigarettes after percutaneous transluminal coronary angioplasty. *American Journal of Cardiology* 61:260-263, 1988.
- GARFINKEL, L. Cancer mortality in nonsmokers: Prospective study by the American Cancer Society. *Journal of the National Cancer Institute* 65(5):1169-1173, November 1980a.
- GARFINKEL, L. Cardiovascular mortality and cigarette smoking. In: Ramstrom, L.M. (ed.) *The Smoking Epidemic. A Matter of Worldwide Concern. Proceedings of The Fourth World Conference on Smoking and Health*. Stockholm: Almqvist and Wiksell International, 1980b, pp. 41-44.
- GARFINKEL, L. Time trends in lung cancer mortality among nonsmokers and a note on passive smoking. *Journal of the National Cancer Institute* 66(6):1061-1066, June 1981.
- GARFINKEL, L. Selection, follow-up and analysis in the American Cancer Society prospective studies. In: Garfinkel, L., Ochs, O., Mushinski, M. (eds.) *Selection, Follow-up, and Analysis in Prospective Studies: A Workshop*, NCI Monograph 67. National Cancer Institute. NIH Publication No. 85-2713, 1985, pp. 49-52.
- GARFINKEL, L., STELLMAN, S.D. Smoking and lung cancer in women. *Cancer Research* 48(23):6951-6955, December 1, 1988.
- GOLDBAUM, G.M., KENDRICK, J.S., HOGELIN, G.C., GENTRY, E.M., BEHAVIORAL RISK FACTORS SURVEYS GROUP. The relative impact of smoking and oral contraceptive use on women in the United States. *Journal of the American Medical Association* 258(10):1339-1342, September 11, 1987.
- HAENZSEL, W., SHIMKIN, M.B., MILLER, H.P., MILMORE, B.K., CONOVER, A.G. *Tobacco Smoking Patterns in the United States*, Public Health Monograph 45. U.S. Public Health Service, 1956.
- HALL, J.R. JR. *Expected Changes in Fire Damages From Reducing Cigarette Ignition Propensity*. Quincy, Massachusetts: National Fire Protection Association, Fire Analysis Division, 1987.

- HAMMOND, E.C. Smoking in relation to physical complaints. *Archives of Environmental Health* 3:146–164, August 1961.
- HAMMOND, E.C. Some preliminary findings on physical complaints from a prospective study of 1,064,004 men and women. *American Journal of Public Health* 54(1):11–23, January 1964a.
- HAMMOND, E.C. Smoking in relation to mortality and morbidity. Findings in the first thirty-four months of follow-up in a prospective study started in 1959. *Journal of the National Cancer Institute* 32(5):1161–1188, May 1964b.
- HAMMOND, E.C. Smoking in relation to the death rates of one million men and women. In: Haenszel, W. (ed.) *Epidemiological Approaches to the Study of Cancer and Other Chronic Diseases*, NCI Monograph 19. U.S. Department of Health, Education, and Welfare, Public Health Service, National Cancer Institute, January 1966, pp. 127–204.
- HAMMOND, E.C. Quantitative relationship between cigarette smoking and death rates. In: Wynder, E.L. (ed.) *Toward a Less Harmful Cigarette*. NCI Monograph 28. U.S. Public Health Service, National Cancer Institute, June 1968, pp. 3–8.
- HAMMOND, E.C. Life expectancy of American men in relation to their smoking habits. *Journal of the National Cancer Institute* 43(4):951–962, October 1969.
- HAMMOND, E.C. Smoking habits and air pollution in relation to lung cancer. In: Lee, D.H.K. (ed.) *Environmental Factors in Respiratory Disease*. New York: Academic Press, 1972.
- HAMMOND, E.C., GARFINKEL, L. Smoking habits of men and women. *Journal of the National Cancer Institute* 27(2):419–442, August 1961.
- HAMMOND, E.C., GARFINKEL, L. Changes in cigarette smoking. *Journal of the National Cancer Institute* 33(1):49–64, July 1964.
- HAMMOND, E.C., GARFINKEL, L. The influence of health on smoking habits. In: Haenszel, W. (ed.) *Epidemiological Approaches to the Study of Cancer and Other Chronic Diseases*, NCI Monograph 19. U.S. Department of Health, Education, and Welfare, Public Health Service, National Cancer Institute, January 1966, pp. 269–285.
- HAMMOND, E.C., GARFINKEL, L. Changes in cigarette smoking 1959–1965. *American Journal of Public Health* 58(1):30–45, January 1968.
- HAMMOND, E.C., GARFINKEL, L. Coronary heart disease, stroke, and aortic aneurysm. Factors in etiology. *Archives of Environmental Health* 19:167–182, August 1969.
- HAMMOND, E.C., GARFINKEL, L. Aspirin and coronary heart disease: Findings of a prospective study. *British Medical Journal* 2:269–271, May 3, 1975.
- HAMMOND, E.C., GARFINKEL, L., SEIDMAN, H., LEW, E.A. “Tar” and nicotine content of cigarette smoke in relation to death rates. *Environmental Research* 12(3):263–274, December 1976.
- HAMMOND, E.C., HORN, D. The relationship between human smoking habits and death rates. A follow-up study of 187,766 men. *Journal of the American Medical Association* 155:1316–1328, August 7, 1954.
- HAMMOND, E.C., SEIDMAN, H. Smoking and cancer in the United States. *Preventive Medicine* 9(2):169–173, March 1980.
- HARRIS, J.E. Cigarette smoking among successive birth cohorts of men and women in the United States during 1900–80. *Journal of the National Cancer Institute* 71(3):473–479, September 1983.
- HAVLIK, R.J., FEINLEIB, M. (eds.) *Proceedings of the Conference on the Decline in Coronary Heart Disease Mortality*. U.S. Department of Health, Education, and Welfare, National Institutes of Health. DHEW Publication No. (NIH) 79-1610, 1979.
- HAWTHORNE, V.M., FRY, J.S. Smoking and health: The association between smoking behaviour, total mortality, and cardiorespiratory disease in west central Scotland. *Journal of Epidemiology and Community Health* 32:260–266, 1978.

- HELLBERG, D., VALENTIN, J., EKLUND, T., STAFFAN, N. Penile cancer: Is there an epidemiological role for smoking and sexual behaviour? *British Medical Journal* 295(6609): 1306–1308, November 21, 1987.
- INTERNATIONAL CLASSIFICATION OF DISEASES. *International Classification of Diseases*, Seventh Revision (ICD-7).
- INTERNATIONAL CLASSIFICATION OF DISEASES. *International Classification of Diseases*, Ninth Revision (ICD-9). U.S. Department of Health and Human Services, Public Health Service, Health Care Financing Administration. DHHS Publication No. (PHS) 80-1260, September 1980.
- KAHN, H.A. The Dorn study of smoking and mortality among U.S. veterans: Report on eight and one-half years of observation. In: Haenszel, W. (ed.) *Epidemiological Approaches to the Study of Cancer and Other Chronic Diseases*, NCI Monograph 19. U.S. Department of Health, Education, and Welfare, Public Health Service, National Cancer Institute, January 1966, pp. 1–125.
- KAUFMAN, D.W., HELMRICH, S.P., ROSENBERG, L., MIETTINEN, O.S., SHAPIRO, S. Nicotine and carbon monoxide content of cigarette smoke and the risk of myocardial infarction in young men. *New England Journal of Medicine* 308(8):409–413, February 24, 1983.
- KINLEN, L.J., ROGOT, E. Leukaemia and smoking habits among United States veterans. *British Medical Journal* 297(6649):657–659, September 10, 1988.
- KLEBBA, A.J. Comparability of mortality statistics for the seventh and eighth revisions of the International Classification of Diseases, United States. *Vital and Health Statistics, Series 2*, No. 66. U.S. Department of Health, Education, and Welfare. DHEW Publication No. (HRA) 76-1340, October 1975.
- KLEBBA, A.J. Mortality from diseases associated with smoking: United States, 1960–77. *Vital and Health Statistics, Series 20*, No. 17. U.S. Department of Health and Human Services. DHHS Publication No. (PHS) 82-1854, January 1982.
- KLEBBA, A.J., SCOTT, J.H. Estimates of selected comparability ratios based on dual coding of 1976 death certificates by the eighth and ninth revisions of the International Classification of Diseases. *Monthly Vital Statistics Report From the National Center for Health Statistics* 28(11):1–19, February 29, 1980.
- KLEBBA, J. Personal communication to J. Harris, June 1988.
- KLEINMAN, J.C., PIERRE, M.B. JR., MADANS, J.H., LAND, G.H., SCHRAMM, W.F. The effects of maternal smoking on fetal and infant mortality. *American Journal of Epidemiology* 127(2):274–282, 1988.
- LAVECCHIA, C., FRANCESCHI, S., DECARLI, A., FASOLI, M., GENTILE, A., TOGNONI, G. Cigarette smoking and the risk of cervical neoplasia. *American Journal of Epidemiology* 123(1):22–29, January 1986.
- LEE, P.N., GARFINKEL, L. Mortality and type of cigarette smoked. *Journal of Epidemiology and Community Health* 35(1):16–22, March 1981.
- LESKO, S.M., ROSENBERG, L., KAUFMAN, D.W., HELMRICH, S.P., MILLER, D.R., STROM, B., SCHOTTENFELD, D., ROSENSHEIN, N.B., KNAPP, R.C., LEWIS, J., SHAPIRO, S. Cigarette smoking and the risk of endometrial cancer. *New England Journal of Medicine* 313(10):593–596, 1985.
- LEVIN, M.L. The occurrence of lung cancer in man. *Acta-Unio Internationalis Contra Cancrum* 9:531–541, 1953.
- LEW, E.A., GARFINKEL, L. Mortality at ages 65 and over in a middle-class population. *Transactions of the Society of Actuaries* 36:257–308, 1984.
- LEW, E.A., GARFINKEL, L. Differences in mortality and longevity by sex, smoking habits and health status. *Transactions of the Society of Actuaries* 39:19–36, 1988.

- LILIENFELD, D.E., GUNDERSON, P.D. The "missing cases" of pleural malignant mesothelioma in Minnesota, 1979-81: Preliminary report. *Public Health Reports* 101(4): 395-399, July-August 1986.
- LILIENFELD, A.M., LILIENFELD, D.E. *Foundations of Epidemiology*, Second Edition. New York: Oxford University Press, 1980.
- LUBIN, J.H., BLOT, W.J., BERRINO, F., FLAMANT, R., GILLIS, C.R., KUNZE, M., SCHMAHL, D., VISCO, D. Modifying risk of developing lung cancer by changing habits of cigarette smoking. *British Medical Journal* 288(6435):1953-1956, June 30, 1984.
- LUBIN, J.H., BOICE, J.D. JR. Estimating radon-induced lung cancer in the U.S. Working paper, 1988.
- MACMAHON, B., PUGH, T.F. *Epidemiology. Principles and Methods*. Boston: Little, Brown and Company, 1970.
- MASSEY, M.M., BOYD, G., MATTSON, M., FEINLEIB, M.R. Inventory of surveys on smoking. *Public Health Reports* 102(4):430-438, July-August 1987.
- MCFARLANE, M.J., FEINSTEIN, A.R., WELLS, C.K. Necropsy evidence of detection bias in the diagnosis of lung cancer. *Archives of Internal Medicine* 146(9):1695-1698, September 1986.
- MCINTOSH, I.D. Smoking and pregnancy: Attributable risks and public health implications. *Canadian Journal of Public Health* 75(2):141-148, March-April 1984.
- MIETTINEN, O.S. Proportion of disease caused or prevented by a given exposure, trait or intervention. *American Journal of Epidemiology* 99(5):325-332, 1974.
- NATIONAL CENTER FOR HEALTH STATISTICS. *Vital Statistics of the United States 1965. Volume 2—Mortality, Part A*. U.S. Department of Health, Education, and Welfare, Public Health Service, National Center for Health Statistics, 1967.
- NATIONAL CENTER FOR HEALTH STATISTICS. Unpublished tabulations, 1985.
- NATIONAL CENTER FOR HEALTH STATISTICS. *Health, United States, 1986*. U.S. Department of Health and Human Services, Public Health Service, National Center for Health Statistics. DHHS Publication No. (PHS) 87-1232, December 1986.
- NATIONAL HEALTH INTERVIEW SURVEY. Unpublished tabulations, 1965.
- NATIONAL HEALTH INTERVIEW SURVEY. Unpublished tabulations, 1985.
- NATIONAL RESEARCH COUNCIL. *Assigned Share for Radiation as a Cause of Cancer. Review of Radioepidemiologic Tables Assigning Probabilities of Causation*. Washington, D.C.: National Academy Press, 1984.
- NATIONAL RESEARCH COUNCIL. *Environmental Tobacco Smoke. Measuring Exposures and Assessing Health Effects*. Washington, D.C.: National Academy Press, 1986.
- NATIONAL RESEARCH COUNCIL. *Health Risks of Radon and Other Internally Deposited Alpha-Emitters. BEIR IV*. Washington, D.C.: National Academy Press, 1988.
- NISCHAN, P., EBELING, K., SCHINDLER, C. Smoking and invasive cervical cancer risk. Results from a case-control study. *American Journal of Epidemiology* 128(1):74-77, July 1988.
- NOVOTNY, T.E., WARNER, K.E., KENDRICK, J.S., REMINGTON, P.L. Smoking by blacks and whites: Socioeconomic and demographic differences. *American Journal of Public Health* 78(9):1187-1189, September 1988.
- OFTEDAL, P., MAGNUS, K., HVINDEN, T. On the probability of radiation being the cause of leukaemia. (Letter.) *British Journal of Radiology* 41:711-712, 1968.
- PEARL, R. Tobacco smoking and longevity. *Science* 87(2253):216-217, March 4, 1938.
- PETO, R. Influence of dose and duration of smoking on lung cancer rates. In: Zaridze, D.G., Peto, R. (eds.) *Tobacco: A Major International Health Hazard*. Lyon, France: International Agency for Research on Cancer, 1986, pp. 23-33.

- RAVENHOLT, R.T. Cigarette smoking: Magnitude of the hazard. (Letter.) *American Journal of Public Health* 54(11):1923-1926, November 1964.
- RAVENHOLT, R.T. Addiction mortality in the United States, 1980: Tobacco, alcohol, and other substances. *Population and Development Review* 10(4):697-724, December 1984.
- RICE, D.P., HODGSON, T.A., SINSHEIMER, P., BROWNER, W., KOPSTEIN, A.N. The economic costs of the health effects of smoking, 1984. *Milbank Quarterly* 64(4):489-547, 1986.
- ROGOT, E. Smoking and mortality among U.S. veterans. *Journal of Chronic Diseases* 27:189-203, 1974.
- ROGOT, E., MURRAY, J.L. Smoking and causes of death among U.S. veterans: 16 years of observation. *Public Health Reports* 95(3):213-222, May-June 1980.
- ROSENBERG, L., KAUFMAN, D.W., HELMRICH, S.P., SHAPIRO, S. The risk of myocardial infarction after quitting smoking in men under 55 years of age. *New England Journal of Medicine* 313(24):1511-1514, December 12, 1985.
- ROSENBERG, L., SCHWINGL, P.J., KAUFMAN, D.W., MILLER, D.R., HELMRICH, S.P., STOLLEY, P.D., SCHOTTENFELD, D., SHAPIRO, S. Breast cancer and cigarette smoking. *New England Journal of Medicine* 310:92-94, January 12, 1984.
- SARACCI, R. The interactions of tobacco smoking and other agents in cancer etiology. *Epidemiologic Reviews* 9:175-193, 1987.
- SCHOENBORN, C.A., BENSON, V. Relationships between smoking and other unhealthy habits: United States, 1985. *Advance Data From Vital and Health Statistics of the National Center for Health Statistics* 154:1-8, May 27, 1988.
- SHULTZ, J.M. SAMMEC. *Smoking-Attributable Mortality, Morbidity, and Economic Costs*. Computer Software and Documentation. Minnesota Department of Health, Center for Non-smoking and Health, 1986.
- SLONE, D., SHAPIRO, S., ROSENBERG, L., KAUFMAN, D., HARTZ, S.C., ROSSI, A.C., STOLLEY, P.D., MIETTINEN, O.S. Relation of cigarette smoking to myocardial infarction in young women. *New England Journal of Medicine* 298(23):1273-1276, June 8, 1978.
- STEIN, Z., KLINE, J., LEVIN, B., SUSSER, M., WARBURTON, D. Epidemiologic studies of environmental exposures in human reproduction. In: Berg, G.G., Maillie, H.D. (eds.) *Measurement of Risks*. New York: Plenum Press, 1981, pp. 163-183.
- STELLMAN, S.D. Cigarette yield and cancer risk: Evidence from case-control and prospective studies. In: Zaridze, D.G., Peto, R. (eds.) *Tobacco: A Major International Health Hazard*. Lyon, France: International Agency for Research on Cancer, 1986a, pp. 197-209.
- STELLMAN, S.D. Influence of cigarette yield on risk of coronary heart disease and chronic obstructive pulmonary disease. In: Zaridze, D.G., Peto, R. (eds.) *Tobacco: A Major International Health Hazard*. Lyon, France: International Agency for Research on Cancer, 1986b, pp. 237-249.
- STELLMAN, S.D., BOFFETTA, P., GARFINKEL, L. Smoking habits of 800,000 American men and women in relation to their occupations. *American Journal of Industrial Medicine* 13:43-58, 1988.
- STELLMAN, S.D., GARFINKEL, L. Smoking habits and tar levels in a new American Cancer Society prospective study of 1.2 million men and women. *Journal of the National Cancer Institute* 76(6):1057-1063, June 1986.
- STERLING, T.D., WEINKAM, J.J. Errors in estimates of smoking-related deaths derived from nonsmoker mortality. *Risk Analysis* 7(4):463-475, 1987.
- TRICHOPOULOS, D., DAY, N.E., KAKLAMANI, E., TZONOOU, A., MUNOZ, N., ZAVITANOS, X., KOUMANTAKI, Y., TRICHOPOULOU, A. Hepatitis B virus, tobacco smoking and ethanol consumption in the etiology of hepatocellular carcinoma. *International Journal of Cancer* 39(1):45-49, January 1987.

- U.S. BUREAU OF THE CENSUS. *Estimates of the Population of the United States, by Age, Sex, and Race: April 1, 1960 to July 1, 1973. Current Population Reports, Series P-25, No. 519.* U.S. Bureau of the Census, April 1974.
- U.S. BUREAU OF THE CENSUS. *Estimates of the Population of the United States, by Age, Sex, and Race: 1980 to 1985. Current Population Reports, Series P-25, No. 985.* U.S. Bureau of the Census, April 1986.
- U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES. *The Health Consequences of Smoking for Women. A Report of the Surgeon General.* U.S. Department of Health and Human Services, Public Health Service, Office of the Assistant Secretary for Health, Office on Smoking and Health, 1980.
- U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES. *The Health Consequences of Smoking. The Changing Cigarette: A Report of the Surgeon General.* U.S. Department of Health and Human Services, Public Health Service, Office of the Assistant Secretary for Health, Office on Smoking and Health. DHHS Publication No. (PHS) 81-50156, 1981.
- U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES. *The Health Consequences of Smoking: Cancer. A Report of the Surgeon General.* U.S. Department of Health and Human Services, Public Health Service, Office on Smoking and Health. DHHS Publication No. (PHS) 82-50179, 1982.
- U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES. *The Health Consequences of Smoking: Cardiovascular Disease. A Report of the Surgeon General.* U.S. Department of Health and Human Services, Public Health Service, Office on Smoking and Health. DHHS Publication No. (PHS) 84-50204, 1983.
- U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES. *The Health Consequences of Smoking: Chronic Obstructive Lung Disease. A Report of the Surgeon General.* U.S. Department of Health and Human Services, Public Health Service, Office on Smoking and Health. DHHS Publication No. (PHS) 84-50205, 1984.
- U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES. *The Health Consequences of Smoking: Cancer and Chronic Lung Disease in the Workplace. A Report of the Surgeon General.* U.S. Department of Health and Human Services, Public Health Service, Office on Smoking and Health. DHHS Publication No. (PHS) 85-50207, 1985.
- U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES. *The Health Consequences of Involuntary Smoking. A Report of the Surgeon General.* U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control. DHHS Publication No. (CDC) 87-8398, 1986.
- U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES. *The Health Consequences of Smoking: Nicotine Addiction. A Report of the Surgeon General, 1988.* U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, Center for Health Promotion and Education, Office on Smoking and Health. DHHS Publication No. (CDC) 88-8406, 1988a.
- U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES. *Health, United States, 1987.* U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Center for Health Statistics. DHHS Publication No. (PHS) 88-1232, March 1988b.
- U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE. *Smoking and Health. A Report of the Surgeon General.* U.S. Department of Health, Education, and Welfare, Public Health Service, Office of the Assistant Secretary for Health, Office on Smoking and Health. DHEW Publication No. (PHS) 79-50066, 1979.
- U.S. OFFICE OF TECHNOLOGY ASSESSMENT. *Smoking-Related Deaths and Financial Costs.* Health Program, Office of Technology Assessment, U.S. Congress, September 1985.

- U.S. PRESIDENT. *Economic Report of the President. Transmitted to the Congress January 1987*. Washington, D.C., 1987.
- U.S. PUBLIC HEALTH SERVICE. *Smoking and Health. Report of the Advisory Committee to the Surgeon General of the Public Health Service*. U.S. Department of Health, Education, and Welfare, Public Health Service, Center for Disease Control. PHS Publication No. 1103, 1964.
- WALTER, S.D. The estimation and interpretation of attributable risk in health research. *Biometrics* 32:829–849, December 1976.
- WHYTE, H.M. The relative importance of the major risk factors in atherosclerotic and other diseases. *Australian and New Zealand Journal of Medicine* 6(5):387–393, October 1976.
- WILCOX, H.B., SCHOENBERG, J.B., MASON, T.J., BILL, J.S., STEMHAGEN, A. Smoking and lung cancer: Risk as function of cigarette tar content. *Preventive Medicine* 17(3):263–272, March 1988.
- WILLETT, W.C., GREEN, A., STAMPFER, M.J., SPEIZER, F.E., COLDITZ, G.A., ROSNER, B., MONSON, R.R., STASON, W., HENNEKENS, C.H. Relative and absolute excess risks of coronary heart disease among women who smoke cigarettes. *New England Journal of Medicine* 317(21):1303–1309, November 19, 1987.
- WILLIAMSON, D.F., FORMAN, M.R., BRADSTOCK, M.K., GENTRY, E., BINKIN, N.J. Behavioral Risk Factor Survey results for cigarette smoking and other health-risk behaviors among U.S. women, 1981–1983. (Abstract.) In: *Epidemic Intelligence Service 35th Annual Conference*. U.S. Department of Health and Human Services, Public Health Service, 1986.
- WYNDER, E.L., STELLMAN, S.D. Impact of long-term filter cigarette usage on lung and larynx cancer risk: A case–control study. *Journal of the National Cancer Institute* 62(3):471–477, March 1979.

CHAPTER 4

TRENDS IN PUBLIC BELIEFS, ATTITUDES, AND OPINIONS ABOUT SMOKING

CONTENTS

Introduction	175
Data Sources	175
Issues in Comparing Surveys	177
Trends in Public Beliefs About the Health Effects of Smoking	179
Overview	179
Is Cigarette Smoking Harmful to Smokers in General?	179
Heavy Versus Light Smoking	181
Tar Yield	183
Duration of Smoking	185
Does Cigarette Smoking Cause:	185
Lung Cancer?	185
Heart Disease?	188
Chronic Obstructive Pulmonary Disease?	188
Other Cancers?	195
What Are the Special Health Risks for Women?	195
Effects of Smoking on Pregnancy Outcome	197
Risk of Cardiovascular Disease Among Smokers	
Who Use Oral Contraceptives	197
Other Health Risks Related to Tobacco Use	200
Involuntary (Passive) Smoking	200
Is Smoking an Addiction?	200
Interaction Between Smoking and Other Exposures	202
Smokeless Tobacco	202
Personal Health Risks for Smokers	202
How Harmful Is Smoking?	204
Absolute Risk	206
Relative Risk	206
Attributable Risk and Smoking-Attributable Mortality	206
Comparative Risk	207
Knowledge Among Adolescents About the Health Risks of Smoking	212
General Health Effects	212
Personalized Risk	215
Comparative Risk	215
Addiction	216
Smokeless Tobacco Use	217
Constituents of Tobacco Smoke	217
Health Benefits of Smoking Cessation	219
Discussion	219
Current Gaps in Public Beliefs About the Health Effects of Smoking ..	219
Factors Interfering With Changes in Knowledge	222
The 1990 Health Objectives for the Nation	223

Trends in Public Attitudes About Smoking and Smokers	224
Involuntary Smoking as an Annoyance	224
Nonsmokers' Rights	224
Actions When Smokers Light Up	227
Opinions of Teenagers	227
<hr/>	
Trends in Public Opinion About Smoking Policies	230
Overview	230
Background	230
Limitations of the Surveys in Assessing Public Opinion	
About Smoking Policies	230
Restrictions on Smoking	230
General	230
Public Places	232
Workplace	232
Airplanes	232
Restaurants	235
Other Places	235
Restrictions on the Sale and Distribution of Cigarettes	235
Complete Ban on Sales	235
Limiting Sales to Minors	235
Banning Free Samples	239
Policies Pertaining to Information and Education	239
Restricting or Prohibiting Tobacco Advertising	239
Warning Labels for Cigarettes	241
Economic Policies	241
Taxation	241
Hiring	241
<hr/>	
Conclusions	244
<hr/>	
Appendix	246
<hr/>	
References	254

Introduction

This Chapter analyzes trends in public beliefs, attitudes, and opinions about smoking. It is divided into three sections. The first describes trends in public beliefs regarding the health effects of smoking, the second describes trends in public attitudes about smokers and smoking, and the third describes trends in public opinion about smoking policies.

At the outset, it is important to define and clarify the important terms used in this Chapter. Terms such as knowledge, awareness, opinions, beliefs, and attitudes have commonsense meanings to the lay person, but more complex meanings to the social scientist. For example, Allport (1935) reviewed many definitions of attitude and constructed his own comprehensive definition: "An attitude is a mental or neural state of readiness, organized through experience, exerting a directive or dynamic influence upon the individual's response to all objects and situations with which it is related." Entire books have been devoted to the science of defining and measuring public attitudes, opinions, and beliefs (e.g., Oskamp 1977).

For sections two and three of this Chapter, which deal with attitudes and opinions, the commonplace understanding of these terms will suffice. For the first section, however, which covers beliefs about health effects, a more careful approach is warranted. This Section generally follows the construct described by Fishbein (1977), which embraces three levels of belief:

1. Level 1 (awareness): A person may believe that "the Surgeon General has determined that cigarette smoking is dangerous to health."
2. Level 2 (general acceptance): A person may believe that "cigarette smoking is dangerous to health."
3. Level 3 (personalized acceptance): A person may believe that "my cigarette smoking is dangerous to my health."

Most of the survey data presented in the first section address Level 2 beliefs. At times, the term public knowledge is used to refer to public beliefs (Level 2 beliefs at the population level). There are few data regarding Level 1 beliefs; consequently, use of the terms awareness and public awareness is generally avoided. Data pertinent to Level 3 beliefs are available from a few surveys in three forms: (1) questions asking whether smoking "is harmful to *your* health"; (2) questions asking whether respondents are "concerned" about the effects of smoking on *their* health; and (3) questions asking whether respondents believe that they are less likely, as likely, or more likely than other people to be adversely affected by smoking. These levels of beliefs are discussed in more depth later in this Chapter.

Data Sources

The information presented in this Chapter is derived from three principal sources:

1. Nationally representative surveys conducted by the U.S. Public Health Service from 1964–87, including the Adult Use of Tobacco Surveys (AUTSs) (1964, 1966, 1970, 1975, 1986) and the National Health Interview Surveys (NHISs) (1985, 1987). The NHIS questions were part of the Health Promotion and Dis-